



MATTHEW D. BARBER, MD, MHS

Department of Obstetrics and Gynecology, The Cleveland Clinic Foundation

KAREN J. MULLEN, MBBS

Lilly Australia, West Ryde, New South Wales

SHERIE A. DOWSETT, PhD, BChD

Eli Lilly Research Laboratories, Indianapolis, IN

LARS VIKTRUP, MD, PhD

Eli Lilly Research Laboratories; and Bowen Research Center, Department of Family Medicine, Indiana University School of Medicine, Indianapolis, IN

The impact of stress urinary incontinence on sexual activity in women

ABSTRACT

In women, stress urinary incontinence is a common problem that may lead to sexual dysfunction. We review the epidemiological data, the pathophysiology, and the risk factors for these two “closet” disorders, how they are related, how we can get patients to talk about them, and how the treatment of stress urinary incontinence may affect sexual dysfunction.

KEY POINTS

More than a third of adult women experience urinary incontinence, with stress urinary incontinence accounting for up to 50% of all cases.

A well-recognized risk factor for stress incontinence is vaginal childbirth: one third of women experience stress incontinence 5 years after their first vaginal delivery.

Leakage of urine during sexual activity is common in women with stress incontinence and may lead to sexual dysfunction.

Leakage appears to occur more frequently during penetration in women with stress urinary incontinence. Women with urge incontinence more often report leakage during orgasm.

MORE THAN A THIRD of adult women experience urinary incontinence,¹ and pure stress urinary incontinence accounts for up to half of all cases.¹ While not a life-threatening disorder, the social^{2,3} and economic⁴ consequences of urinary incontinence may be profound. Nevertheless, it remains a “closet” problem, as an appreciable number of women do not seek treatment, even when their symptoms cause significant distress and interfere with daily activities.^{2,5,6}

In some cases, wearing pads or modifying daily activities helps limit embarrassing situations.^{2,3,7} When leakage occurs during sexual activity, however, the disorder may be more difficult to cope with, and the psychological impact⁸ and the strain on a relationship⁹ may be significant.

Defining what role stress urinary incontinence plays in sexual dysfunction is challenging, due in part to the complex etiology of sexual dysfunction and to the taboo nature of both disorders. The aim of this article is to review the epidemiologic data and discuss our current understanding of the relationship between these two disorders and the challenges faced in addressing this topic with patients.

WHAT IS STRESS URINARY INCONTINENCE, AND HOW COMMON IS IT?

Stress urinary incontinence is defined as the complaint of involuntary leakage of urine on effort or exertion, or on sneezing or coughing,¹⁰ usually resulting from urethral or bladder neck hypermobility or urethral sphincter insufficiency. A diagnosis is provisionally based on the patient’s history and signs of

Dr. Dowsett, Mullen, and Viktrup are full-time employees of Eli Lilly and Company and are stockholders in the Lilly Company. Dr. Barber received a \$2,000 honorarium for his role as an author of this manuscript.

This paper discusses therapies that are experimental or are not approved by the US Food and Drug Administration for the use under discussion.

TABLE 1

Potential risk factors for stress urinary incontinence

Age
Pregnancy
Vaginal childbirth
Obesity
Constipation
Smoking
Chronic respiratory disease
Use of alpha-blockers
Pelvic surgery

leakage during the stress test, and is confirmed through urodynamic evaluation.¹⁰

In contrast, urge incontinence is the complaint of involuntary leakage accompanied or immediately preceded by a feeling of urgency.¹⁰ Urge incontinence usually results from overactivity of the detrusor muscle that may (in the case of motor urgency) or may not (in the case of sensory urgency) be confirmed during urodynamic testing.

Prevalence increases with age

The true prevalence of stress urinary incontinence is open to conjecture, as it is for urinary incontinence in general, since sampling techniques, case definitions, and the questions asked of study participants differ dramatically between studies. Moreover, underreporting is to be expected, given the embarrassment associated with the problem. Yet the problem is more prevalent with increasing age. A recent summary¹ estimates the prevalence of urinary incontinence in general to be from 20% to 50%, with a broad peak during middle age (30% to 40%) and a steady increase in the elderly (30% to 50%).

Stress incontinence is more prevalent than other types

Pure stress incontinence accounts for half of all cases of urinary incontinence in women, but a component of stress incontinence is seen in three fourths of all cases of urinary incontinence.¹ A meta-analysis by Hampel et al¹¹ revealed similar findings: In data published from 1954 to 1995, 10% to 50% of women

reported symptoms of urinary incontinence, with pure stress incontinence accounting for 49% of all incontinence cases, and a stress component evident in a further 29%.¹¹

In a recent nationwide US survey of 24,581 community-dwelling women, 9,002 (37%) reported incontinence symptoms within the past 30 days.¹² Of these, 41% reported stress incontinence only, with a further 45% reporting symptoms of both stress and urge incontinence, thus totaling more than 85% of women with urinary incontinence with a stress component.

Risk factors for stress incontinence

In addition to age,¹ another well-recognized risk factor for stress incontinence is vaginal childbirth, with one third of women experiencing stress incontinence 5 years after their first delivery,¹³ presumably due to neuromuscular damage to the urethral sphincter or pelvic floor, or to injury to the supporting connective tissue. Other risk factors have not been as vigorously studied, but may include obesity,^{14,15} constipation,¹⁴ smoking,¹⁶ chronic obstructive lung disease,¹⁷ use of alpha-blockers,¹⁸ and pelvic surgery^{19,20} (TABLE 1).

■ WHAT IS FEMALE SEXUAL DYSFUNCTION, AND HOW COMMON IS IT?

The definition of sexual dysfunction has evolved since the early 1990s as has our understanding of the normal cycle of female sexual response. This cycle, initially defined by Masters and Johnson,²¹ was later divided by Kaplan²² into three phases—desire, arousal, and orgasm. This forms the basis for the current definition of female sexual dysfunction, as proposed by American Foundation of Urologic Disease in 1998.²³ Using this classification, sexual dysfunction encompasses:

- Hypoactive sexual desire and sexual aversion disorders (deficiency of sexual thoughts or receptivity to sexual activity that causes personal distress)
- Sexual arousal disorder (inability to attain or maintain sufficient sexual excitement that causes personal distress)
- Orgasmic disorder
- Sexual pain disorders, which include dyspareunia (genital pain associated with sexual

1/3 of women have some stress urinary incontinence 5 years after their first vaginal birth



intercourse) and vaginismus (involuntary spasm of the musculature of the outer third of the vagina interfering with penetration).

Note that the diagnosis requires that symptoms be severe enough to cause personal distress.²³

Prevalence of sexual dysfunction

Establishing an estimate for the prevalence of sexual dysfunction in women in the general population is plagued with challenges similar to those for urinary incontinence. For example, the case definition is not standardized, and underreporting is surely to be expected, given social taboos associated with it. Ideally, evaluation of sexual dysfunction should encompass a physical examination, a psychological and psychosocial assessment, and, in some cases, an endocrine evaluation. To limit the embarrassment of answering sensitive questions on sexual activity openly, the assessment should use a questionnaire that can be filled out at home, since self-reporting tools have been shown to provide a more valid measurement of sexual function.^{24,25}

By far the most commonly cited prevalence rate of sexual dysfunction in US women is 43%.²⁶ This finding was from a study in 2000 of 1,749 women ages 18 to 59 participating in the National Health and Social Life Survey. Importantly, sexual dysfunction in this case was not based on clinical diagnosis, but was defined only as a report of lack of interest in sex, inability to achieve orgasm, pain during sex, sex that was not pleasurable, anxiety about performance, or trouble lubricating. It is not clear how many of these women experienced personal distress, as required for diagnosis.²³

Causes of sexual dysfunction

The etiology of female sexual dysfunction is complex and encompasses biological, psychological, and interpersonal factors²³ (TABLE 2). Biological factors may include reduced vascular flow to the genital tissues, or injury or disease of the central or peripheral nervous system. Psychological factors include lack of emotional intimacy²⁷ and depression.²⁸ In addition, drugs used to treat depression can also adversely affect sexual function.²⁹

Other risk factors for sexual dysfunction in women include a history of traumatic sexu-

TABLE 2

Factors associated with female sexual dysfunction

- Urinary incontinence
- Age
- Lack of emotional intimacy
- Depression
- Antidepressant drugs
- History of a traumatic sexual event
- Diabetes
- Hypertension
- Pelvic inflammatory disease
- Pelvic trauma
- Menopause

al events,²⁶ diabetes,³⁰ hypertension,³¹ pelvic inflammatory disease,³² or pelvic trauma.³³ Menopause and the ensuing lowered estrogen levels may also result in sexual dysfunction,²¹ for example, as a consequence of decreased lubrication. Conversely, the prevalence of sexual problems, except for inadequate lubrication, generally decreases with age.²⁶

■ DOES LEAKAGE OCCUR DURING SEXUAL ACTIVITY?

Studies suggest urinary leakage occurs during sexual activity in 11% to 60% of women with stress urinary incontinence.^{3,8,34-45} This large range likely reflects methodological and population differences between studies and small sample sizes.

Study populations have consisted primarily of women attending clinics specifically focusing on urologic concerns, although studies of general populations have also been conducted. In a study by Lam et al,³ a self-administered questionnaire was mailed to a random sample of women living in Aarhus, Denmark. Overall, 12% of 441 women with stress urinary incontinence reported leakage during sexual intercourse. Nygaard and Milburn⁴⁶ provided mail-in questionnaires to all non-pregnant women presenting for routine annual gynecologic examination in selected clinics in Iowa over a 1-month period. Of the 224 who returned the questionnaire, 77% reported urinary leakage to some degree (“mild” or

A diagnosis of sexual dysfunction requires that symptoms cause personal distress

TABLE 3

Measures of the impact of urinary incontinence on quality of life and sexual function in women

Generic measures of quality of life

Medical Outcomes Study Short Form (SF-36)⁵⁰
EuroQol/EQ-5D⁵¹
Sickness Impact Profile (SIP)⁵²
Nottingham Health Profile Questionnaire⁵³

Measures of quality of life specific to urinary incontinence

Urogenital Distress Inventory (UDI)⁵⁴
Bristol Female Lower Urinary Tract Symptoms Questionnaire (B-FLUTS)⁵⁵
Urinary Incontinence Quality of Life Instrument (I-QoL)⁵⁶
Incontinence Impact Questionnaire (IIQ)⁵⁷
Kings Health Questionnaire⁵⁸

Measures of quality of life specific to sexual function

Golombok-Rust Inventory of sexual satisfaction⁶⁰
Psychosocial Adjustment to Illness Scale (PAIS)⁶¹
Brief Index of Sexual Functioning for Women (BISF-W)⁶²
Female Sexual Function Index (FSFI)⁶³
Derogatis Interview for Sexual Functioning (DISF/DISF-SR)⁶⁴
Simple Sexual Function Questionnaire⁶⁵

Measures of sexual function specific to urinary incontinence

B-FLUTS (Sex) Questionnaire⁵⁵
I-QoL⁵⁶
IIQ⁵⁷
Kings Health Questionnaire⁵⁸
Pelvic Organ Prolapse/Urinary Incontinence Sexual Function Questionnaire (PISQ)⁴⁹
PISQ Short Form (PISQ-12)⁵⁹

The etiology of female sexual dysfunction is complex

“moderate”), with 12% of those with mild incontinence and 24% of those with moderate incontinence reporting urine loss during sexual activity. In this study, stress urinary incontinence was not specifically discussed, although four of the seven nonsexual questions focused on urinary incontinence with physical exertion. Interestingly, in this study 2 (4%) of the 52 women who did not report incontinent episodes based on the nonsexual questioning did report incontinence during sexual activity.

■ AT WHAT STAGE OF SEXUAL ACTIVITY DOES LEAKAGE OCCUR?

Several survey studies have included questions designed to determine at what point during sexual activity leakage is likely to occur. In a case-control study by Hilton,³⁶ 48 (23%) of the women with stress incontinence experienced incontinence during intercourse. Of

these women, 18% reported incontinence during penetration and 5% during orgasm. These results are supported by those of Clark and Romm,³⁸ who concluded that leakage was more likely to occur during penetration than during arousal, orgasm, or resolution. In a later study of 57 incontinent women who reported leakage during sexual activity, 14 were diagnosed with pure stress incontinence through questioning, and 12 (86%) of these reported urine loss during deep penetration.⁴⁰ Urine leakage during other sexual activities was not described in this study. Berglund and Fugl-Meyer⁴¹ found that of 44 women with stress urinary incontinence who were referred for surgery, 12 (27%) had reported leakage related to the penetration movements during intercourse (prior to surgery), and a further 13 (30%) were unsure if they had experienced leakage during intercourse. While 39% of these 44 women reported orgasmic dysfunction



tion, leakage as a direct cause of orgasmic dysfunction (that is, fear of leakage that led to distraction) was volunteered by only 2 women. In another study,⁴⁶ women undergoing a routine gynecologic examination reported that leakage was more common during orgasm than penetration, although numbers reporting these specific concerns were small (6 vs 2 women) and the diagnosis of urinary incontinence was not specific to stress incontinence.⁴⁶

Based on the limited data, leakage appears to occur more frequently during penetration in women with stress urinary incontinence,^{36,38} but women with urge incontinence more often report leakage during orgasm.^{36,38}

■ WHAT IS THE IMPACT OF LEAKAGE ON SEXUAL ACTIVITY?

One can readily appreciate that a woman who has experienced urinary leakage during sexual activity is likely to be apprehensive of similar episodes in the future. In fact, while urinary leakage during sexual activity generates immediate embarrassment, the long-term psychological impact is likely to be of greater significance.⁸ Women with stress incontinence report a number of factors that contribute to a decreased interest in sexual activity: loss of spontaneity, the need for washing or for disposal of wet pads, the need for separate beds, a general feeling of unattractiveness from wearing pads in bed, and concerns over odor.^{34,38,47}

Sexual expression and behavior are also likely to be affected in the broader sense, that is, in the outward expression of femininity; for example, in terms of the appropriate choice of clothing in case of leakage.⁴⁸ In essence, a woman's perception of her own sexuality is jeopardized, leading to significant inhibition. Further stigma may ensue based on her partner's reaction.

How to measure the impact

To evaluate the impact of urinary incontinence on quality of life, a number of questionnaires have been developed and validated (TABLE 3).^{42,49–66} These measure the severity of urinary symptoms and their impact on activities of daily living, physical activities, social interactions, personal relationships, and self-perception; some specifically address the

impact on sexual function.

The King's Health Questionnaire, the Incontinence Quality of Life Instrument (I-QoL), and the Incontinence Impact Questionnaire (IIQ) contain one or two questions to evaluate the impact on the enjoyment of sex, on sex life, or on sexual relations. The Bristol Female Lower Urinary Tract Symptoms (B-FLUTS) questionnaire⁴² includes four questions relating to sexual function:

- Pain or discomfort due to a dry vagina
- Sex life spoiled by urinary symptoms
- Pain on sexual intercourse
- Leakage on intercourse.

This questionnaire has been used to evaluate the effectiveness of pelvic floor muscle training on quality of life and sexual health⁴³ and to determine the effect of urinary incontinence on sexual activity in a community-based population (see below).⁶⁶

The Pelvic Organ Prolapse/Urinary Incontinence Sexual Function Questionnaire (PISQ) was developed and validated in women with symptoms of either pelvic organ prolapse or urinary incontinence.⁴⁹ This questionnaire focuses specifically on urinary incontinence and sexual health and comprises behavioral-emotive, physical, and partner-related domains. A short form of the questionnaire, the PISQ-12, is also available.⁵⁹ Currently, the PISQ is the only condition-specific sexual function questionnaire dedicated to women with urinary incontinence. Questionnaires to specifically assess sexual function irrespective of continence status may also be of some value to measure the impact of stress urinary incontinence on sexual function (TABLE 3).

■ DOES TREATING INCONTINENCE RESTORE SEXUAL SATISFACTION?

Pelvic floor muscle training

Pelvic floor exercises have been shown to reduce symptoms of urinary leakage in women with stress urinary incontinence,^{67,68} improve quality of life in general⁴³ and, more specifically, improve overall sex life.⁴³ In a study by Bo et al,⁴³ urinary incontinence with intercourse was reduced by nearly 50% in women who performed pelvic floor exercises vs less than 10% in the control group, who received no intervention.⁴³

Many women do not seek treatment for incontinence

Regrettably, studies have generally been small and lacked an appropriate control, and outcomes measures have been inconsistent across trials. Despite this, pelvic floor exercises pose little potential for harm and are of minimal cost; therefore, they should be considered a first-line treatment for stress urinary incontinence. For women with both stress incontinence and sexual dysfunction, pelvic floor exercises present a unique option that may be effective for both conditions.

Surgery

Surgical treatment of stress urinary incontinence is recommended when more conservative treatment has failed.⁶⁹ In a US study using a mailed survey, 4% of 24,000 community-dwelling women reported a history of continence surgery.⁷ Unfortunately, assessing the efficacy of various surgical methods is problematic due to inadequate case definitions, use of unvalidated outcome tools, inconsistent outcome measures, and concerns over the inability to generalize the results.⁷⁰ In spite of this, two systematic reviews have concluded that retropubic colposuspension (the Burch and Marshall-Marchetti-Krantz procedures) and placement of a suburethral sling are the most effective surgical procedures for treatment of stress urinary incontinence.^{69,70}

Minimally invasive outpatient procedures introduced more recently, such as the tension-free vaginal tape or “TVT” procedure, have a reported efficacy similar to that for the Burch colposuspension procedure at 2 years.⁷¹

Surgery is not without risk, even though it is effective in many cases. As many as 23% of patients who undergo surgery for stress incontinence develop postoperative voiding dysfunction, and 10% develop new-onset detrusor overactivity.⁷² In addition, 3% to 9% of patients treated surgically have been reported to have urinary tract injury.^{73,74}

Cure rates with surgery range from 51% to 91%, depending on the analytic methods, the length of follow-up, and the definition of cure.^{69,71} Several studies have demonstrated that surgery for stress incontinence can result in an improvement in sexual function,^{41,44,75,76} although there is also potential for reduced sexual enjoyment in a subset of surgically treated women.

Sexual dissatisfaction after surgery is often due to dyspareunia,^{76,77} but orgasmic dysfunction has also been reported to play a role.⁴¹ In addition, feelings of anxiousness and distress from worrying about potential urinary leakage during sexual activity may not be resolved once stress incontinence is treated. This, together with unrealistic patient expectations,⁴¹ may also explain the sexual dissatisfaction after surgery in some women.

Drug therapy

No drug has yet been approved worldwide for treating stress urinary incontinence, although alpha-adrenergic receptor agonists and tricyclic antidepressants have been used off-label.⁷⁸ Estrogen preparations, both systemic and topical, have also been used, but several randomized trials have not found them to be effective.⁷⁹ There is some evidence, however, that estrogen may be useful for treating urge incontinence and other irritative voiding symptoms.⁸⁰ Furthermore, topical estrogen therapy can be effective for treating vulvovaginal atrophy and vaginal dryness,⁸¹ which can contribute to sexual dysfunction in postmenopausal women.

Educate patients about what to expect

While stress incontinence and sexual dysfunction often coexist, stress incontinence may not be the prime cause of sexual dysfunction in many women. To successfully address a woman's individual needs, one should consider sexual dysfunction in the larger context of all factors that could contribute to it. It is also important that the patient's expectations be reasonable: ie, she should know that, while treatment may reduce symptoms during sexual activity, no cure is guaranteed.

■ IMPLICATIONS FOR PRIMARY CARE PHYSICIANS

Both stress urinary incontinence and sexual dysfunction affect many women and significantly worsen quality of life. Yet women are often reluctant to volunteer information about their symptoms because of the stigma attached to these problems.

Given the psychosocial consequences of these conditions and the increasing availabili-

Pelvic floor exercises are the first-line treatment for stress urinary incontinence



ty of safe and effective treatments, primary care physicians should be prepared to ask about these disorders. One way is to add simple questions such as “Do you leak urine during physical activity or when you cough or sneeze?” and “Do you have any difficulties with your sex life?” to the review of systems during routine health screenings and annual gynecologic examinations. The urinary incontinence and sexual function questionnaires in TABLE 3 are also useful for this.

Many women with stress urinary incontinence may leak urine during sexual intercourse, with potential for immediate embarrassment and decreased desire and avoidance of sexual activity in the longer term. Quality research is needed to further characterize the relationship between these two conditions and to further elucidate the impact, whether positive or negative, of treatment of stress incontinence on the sexual function of women.

REFERENCES

- Hunnskaar S, Arnold EP, Burgio K, Diokno AC, Herzog AR, Mallett VT. Epidemiology and natural history of urinary incontinence. *Int Urogynecol J Pelvic Floor Dysfunct* 2000; 11:301-319.
- Fultz H, Burgio K, Diokno AC, Kinchen KH, Obenchain R, Bump RC. Burden of stress urinary incontinence for community-dwelling women. *Am J Obstet Gynecol* 2003; 189:1275-1282.
- Lam GW, Foldspang A, Elving LB, Mommsen S. Social context, social abstinence, and problem recognition correlated with adult female urinary incontinence. *Dan Med Bull* 1992; 39:565-570.
- Wagner TH, Hu TW. Economic costs of urinary incontinence in 1995. *Urology* 1998; 51:355-361.
- Burgio KL, Ives DG, Locher JL, Arena VC, Kuller LH. Treatment seeking for urinary incontinence in older adults. *J Am Geriatr Soc* 1994; 42:208-212.
- Diokno AC. Epidemiology and psychosocial aspects of incontinence. *Urol Clin North Am* 1995; 22:481-485.
- Diokno AC, Burgio K, Fultz H, Kinchen KH, Obenchain R, Bump RC. Medical and self-care practices reported by women experiencing urinary incontinence. *Am J Manag Care*. In press 2003.
- Kelleher C, Cardozo L, Khullar V, Wise B, Cutner A. The impact of urinary incontinence on sexual function. *J Sexual Health*, Winter 1993:186-191.
- Yip SK, Chan A, Pang S, et al. The impact of urodynamic stress incontinence and detrusor overactivity on marital relationship and sexual function. *Am J Obstet Gynecol* 2003; 188:1244-1248.
- Abrams P, Cardozo L, Fall M, et al. The standardisation of terminology of lower urinary tract function: report from the Standardisation Sub-committee of the International Continence Society. *Neurourol Urodyn* 2002; 21:167-178.
- Hampel C, Wienhold D, Benken N, Eggersmann C, Thuroff JW. Prevalence and natural history of female incontinence. *Eur Urol* 1997; 32(suppl 12):3-12.
- Diokno AC, Burgio K, Fultz H, Kinchen KH, Obenchain R, Bump RC. Prevalence and outcomes of continence surgery in community dwelling women. *J Urology* 2003; 170:507-511.
- Viktrup L, Lose G. The risk of stress incontinence 5 years after first delivery. *Am J Obstet Gynecol* 2001; 185:82-87.
- Moller LA, Lose G, Jorgensen T. Risk factors for lower urinary tract symptoms in women 40 to 60 years of age. *Obstet Gynecol* 2000; 96:446-451.
- Hannestad YS, Rortveit G, Daltveit AK, Hunnskaar S. Are smoking and other lifestyle factors associated with female urinary incontinence? The Norwegian EPINCONT Study. *BJOG* 2003; 110:247-254.
- Bump RC, McClish DK. Cigarette smoking and urinary incontinence in women. *Am J Obstet Gynecol* 1992; 167:1213-1218.
- Brown JS, Seeley DG, Fong J, Black DM, Ensrud KE, Grady D. Urinary incontinence in older women: who is at risk? Study of Osteoporotic Fractures Research Group. *Obstet Gynecol* 1996; 87:715-721.
- Menefee SA, Chesson R, Wall LL. Stress urinary incontinence due to prescription medications: alpha-blockers and angiotensin-converting enzyme inhibitors. *Obstet Gynecol* 1998; 91:853-854.
- Mommsen S, Foldspang A, Elving L, Lam GW. Association between urinary incontinence in women and a previous history of surgery. *Br J Urol* 1993; 72:30-37.
- Reid G. C., DeLancey JO, Hopkins MP, Roberts JA, Morley GW. Urinary incontinence following radical vulvectomy. *Obstet Gynecol* 1990; 75:852-858.
- Masters WH, Johnson VE. *Human Sexual Response*. St. Louis, MO: Little, Brown, 1966.
- Kaplan HS. *The New Sex Therapy: Active Treatment of Sexual Dysfunctions*. New York: Brunner and Mazel, 1974.
- Basson R, Berman J, Burnett A, et al. Report of the international consensus development conference on female sexual dysfunction: definitions and classifications. *J Urol* 2000; 163:888-893.
- Rosen RC, Beck JG. *Patterns of Sexual Arousal: Psychophysiological Processes and Clinical Applications*. New York: Guilford Press, 1988.
- Tourangeau R, Smith TW. Asking sensitive questions: the impact of data collection mode, question format and question context. *Public Opinion Quarterly* 2003; 60:275-304.
- Laumann EO, Paik A, Rosen RC. Sexual dysfunction in the United States: prevalence and predictors. *JAMA* 1999; 281:537-544.
- Basson R. Human sex-response cycles. *J Sex Marital Ther* 2001; 27:33-43.
- Casper RC, Redmond DE Jr, Katz MM, Schaffer CB, Davis JM, Koslow SH. Somatic symptoms in primary affective disorder. Presence and relationship to the classification of depression. *Arch Gen Psychiatry* 1985; 42:1098-1104.
- Rosen RC, Lane RM, Menza M. Effects of SSRIs on sexual function: a critical review. *J Clin Psychopharmacol* 1999; 19:67-85.
- Schreiner-Engel P, Schiavi RC, Vietorisz D, Smith H. The differential impact of diabetes type on female sexuality. *J Psychosom Res* 1987; 31:23-33.
- Duncan LE, Lewis C, Jenkins P, Pearson TA. Does hypertension and its pharmacotherapy affect the quality of sexual function in women? *Am J Hypertens* 2000; 13:640-647.
- Heisterberg L. Factors influencing spontaneous abortion, dyspareunia, dysmenorrhea, and pelvic pain. *Obstet Gynecol* 1993; 81:594-597.
- Copeland CE, Bosse MJ, McCarthy ML, et al. Effect of trauma and pelvic fracture on female genitourinary, sexual, and reproductive function. *J Orthop Trauma* 1997; 11:73-81.
- Sutherst JR. Sexual dysfunction and urinary incontinence. *Br J Obstet Gynaecol* 1979; 86:387-388.
- Sutherst JR, Brown M. Sexual dysfunction associated with urinary incontinence. *Urol Int* 1980; 35:414-416.
- Hilton P. Urinary incontinence during sexual intercourse: a common, but rarely volunteered, symptom. *Br J Obstet Gynaecol* 1988; 95:377-381.
- Walters MD, Taylor S, Schoenfeld LS. Psychosexual study of women with detrusor instability. *Obstet Gynecol* 1990; 75:22-26.
- Clark A, Romm J. Effect of urinary incontinence on sexual activity in women. *J Reprod Med* 1993; 38:679-683.
- Field SM, Hilton P. The prevalence of sexual problems in women attending for urodynamic investigation. *Int Urogynecol J Pelvic Floor*



- Dysfunct 1993; 4:212–215.
40. **Vierhout ME, Gianotten WL.** Mechanisms of urine loss during sexual activity. *Eur J Obstet Gynecol Reprod Biol* 1993; 52:45–47.
 41. **Berglund AL, Fugl-Meyer KS.** Some sexological characteristics of stress incontinent women. *Scand J Urol Nephrol* 1996; 30:207–212.
 42. **Jackson S, Shepherd A, Brookes S, Abrams P.** The effect of oestrogen supplementation on post-menopausal urinary stress incontinence: a double-blind placebo-controlled trial. *Br J Obstet Gynaecol* 1999; 106:711–718.
 43. **Bo K, Talseth T, Vinsnes A.** Randomized controlled trial on the effect of pelvic floor muscle training on quality of life and sexual problems in genuine stress incontinent women. *Acta Obstet Gynecol Scand* 2000; 79:598–603.
 44. **Lemack GE, Zimmern PE.** Sexual function after vaginal surgery for stress incontinence: results of a mailed questionnaire. *Urology* 2000; 56:223–227.
 45. **Barber MD, Visco AG, Wyman JF, Fantl JA, Bump RC.** Continence Program for Women Research Group. Sexual function in women with urinary incontinence and pelvic organ prolapse. *Obstet Gynecol* 2002; 99:281–289.
 46. **Nygaard I, Milburn A.** Urinary incontinence during sexual activity: prevalence in a gynecologic practice. *J Womens Health* 1995; 4:83–86.
 47. **Norton C.** The effects of urinary incontinence in women. *Int Rehabil Med* 1982; 4:9–14.
 48. **Roe B, May C.** Incontinence and sexuality: findings from a qualitative perspective. *J Adv Nurs* 1999; 30:573–579.
 49. **Rogers RG, Kammerer-Doak D, Villarreal A, Coates K, Qualls C.** A new instrument to measure sexual function in women with urinary incontinence or pelvic organ prolapse. *Am J Obstet Gynecol* 2001; 184:552–558.
 50. **Ware JE Jr, Sherbourne CD.** The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992; 30:473–483.
 51. **EuroQol—a new facility for the measurement of health-related quality of life.** The EuroQol Group. *Health Policy* 1990; 16:199–208.
 52. **Hunnskaar S, Vinsnes A.** The quality of life in women with urinary incontinence as measured by the sickness impact profile. *J Am Geriatr Soc* 1991; 39:378–382.
 53. **Grimby A, Milsom I, Molander U, Wiklund I, Ekelund P.** The influence of urinary incontinence on the quality of life of elderly women. *Age Ageing* 1993; 22:82–89.
 54. **Shumaker SA, Wyman JF, Uebersax JS, McClish D, Fantl JA.** Health-related quality of life measures for women with urinary incontinence: the Incontinence Impact Questionnaire and the Urogenital Distress Inventory. Continence Program in Women (CPW) Research Group. *Qual Life Res* 1994; 3:291–306.
 55. **Jackson S, Donovan J, Brookes S, Eckford S, Swithinbank L, Abrams P.** The Bristol Female Lower Urinary Tract Symptoms questionnaire: development and psychometric testing. *Br J Urol* 1996; 77:805–812.
 56. **Wagner TH, Patrick DL, Bavendam TG, Martin ML, Buesching DP.** Quality of life of persons with urinary incontinence: development of a new measure. *Urology* 1996; 47:67–71.
 57. **Wyman JF, Harkins SW, Choi SC, Taylor JR, Fantl JA.** Psychosocial impact of urinary incontinence in women. *Obstet Gynecol* 1987; 70:378–381.
 58. **Kelleher CJ, Cardozo LD, Khullar V, Salvatore S.** A new questionnaire to assess the quality of life of urinary incontinent women. *Br J Obstet Gynaecol* 1997; 104:1374–1379.
 59. **Rogers RG, Coates KW, Kammerer-Doak D, Khalsa S, Qualls C.** A short form of the Pelvic Organ Prolapse/Urinary Incontinence Sexual Questionnaire (PISQ-12). *Int Urogynecol J Pelvic Floor Dysfunct* 2003; 14:164–168.
 60. **Rust J, Golombok S.** The Golombok-Rust Inventory of Sexual Satisfaction (GRISS). *Br J Clin Psychol* 1985; 24:63–64.
 61. **Derogatis LR.** The psychosocial adjustment to illness scale (PAIS). *J Psychosom Res* 1986; 30:77–91.
 62. **Taylor JF, Rosen RC, Leiblum SR.** Self-report assessment of female sexual function: psychometric evaluation of the Brief Index of Sexual Functioning for Women. *Arch Sex Behav* 1994; 23:627–643.
 63. **Rosen R, Brown C, Heiman J, et al.** The Female Sexual Function Index (FSFI): a multidimensional self-report instrument for the assessment of female sexual function. *J Sex Marital Ther* 2000; 26:191–208.
 64. **Derogatis LR.** The Derogatis Interview for Sexual Functioning (DISF/DISF-SR): an introductory report. *J Sex Marital Ther* 1997; 23:291–304.
 65. **Plouffe L Jr.** Screening for sexual problems through a simple questionnaire. *Am J Obstet Gynecol* 1985; 151:166–169.
 66. **Temml C, Haidinger G, Schmidbauer J, Schatzl G, Madersbacher S.** Urinary incontinence in both sexes: prevalence rates and impact on quality of life and sexual life. *Neurourol Urodyn* 2000; 19:259–271.
 67. **Bo K, Talseth T, Holme I.** Single blind, randomised controlled trial of pelvic floor exercises, electrical stimulation, vaginal cones, and no treatment in management of genuine stress incontinence in women. *BMJ* 1999; 318:487–493.
 68. **Hay-Smith EJ, Bo Berghmans LC, Hendriks HJ, de Bie RA, van Waalwijk van Doorn ES.** Pelvic floor muscle training for urinary incontinence in women. *Cochrane Database Syst Rev.* 2001; (1):CD001407.
 69. **Leach GE, Dmochowski RR, Appell RA, et al.** Female Stress Urinary Incontinence Clinical Guidelines Panel summary report on surgical management of female stress urinary incontinence. The American Urological Association. *J Urol* 1997; 158:875–880.
 70. **Black NA, Downs SH.** The effectiveness of surgery for stress incontinence in women: a systematic review. *Br J Urol* 1996; 78:497–510.
 71. **Ward KL, Hilton P.** UK and Ireland TVT Trial Group. A prospective multicenter randomized trial of tension-free vaginal tape and colposuspension for primary urodynamic stress incontinence: two-year follow-up. *Am J Obstet Gynecol* 2004; 190:324–331.
 72. **Smith T, Daneshgari FDR.** Surgical treatment of incontinence in women. In: Abrams P, Cardozo L, Khoury S, Wein A, editors. 2nd International consultation on incontinence. Plymouth, UK: Health Publications, 2002.
 73. **Tulikangas PK, Weber AM, Larive AB, Walters MD.** Intraoperative cystoscopy in conjunction with anti-incontinence surgery. *Obstet Gynecol* 2000; 95:794–796.
 74. **Ward K, Hilton P, United Kingdom and Ireland Tension-free Vaginal Tape Trial Group.** Prospective multicentre randomised trial of tension-free vaginal tape and colposuspension as primary treatment for stress incontinence. *BMJ* 2002; 325:67.
 75. **Black NA, Bowling A, Griffiths JM, Pope C, Abel PD.** Impact of surgery for stress incontinence on the social lives of women. *Br J Obstet Gynaecol* 1998; 105:605–612.
 76. **Haase P, Skibsted L.** Influence of operations for stress incontinence and/or genital descensus on sexual life. *Acta Obstet Gynecol Scand* 1988; 67:659–661.
 77. **Weber AM, Walters MD, Piedmonte MR.** Sexual function and vaginal anatomy in women before and after surgery for pelvic organ prolapse and urinary incontinence. *Am J Obstet Gynecol* 2000; 182:1610–1615.
 78. **Viktrup L, Bump RC.** Pharmacological agents used for the treatment of stress urinary incontinence in women. *Curr Med Res Opin* 2003; 19:485–490.
 79. **Al Badr A, Ross S, Soroka D, Drutz HP.** What is the available evidence for hormone replacement therapy in women with stress urinary incontinence?. *J Obstet Gynaecol Can* 2003; 25:567–574.
 80. **Cardozo L, Lose G, McClish D, Versi E.** Estrogen treatment for symptoms of an overactive bladder: Results of a meta-analysis. *Int Urogynecol J Pelvic Floor Dysfunct* 2001; 12(suppl 3):S43.
 81. **Blakeman PJ, Hilton P, Bulmer JN.** Cellular proliferation in the female lower urinary tract with reference to oestrogen status. *BJOG* 2001; 108:813–816.

ADDRESS: Lars Viktrup, MD, PhD, Eli Lilly and Company, Faris II, Drop Code 6112, Indianapolis, IN 46285; e-mail Viktrup_lars@lilly.com.